

THE FIRST RETURNING SPEED AND THE LAST EXIT SPEED OF A TYPE OF MARKOV CHAIN

HUIZENG ZHANG*, MINZHI ZHAO**, AND LEI WANG***

ABSTRACT. Let $\{X_n\}$ be a Markov chain with transition probability $p_{ij} = a_{j-(i-1)+}, \forall i, j \geq 0$, where $a_j = 0$ provided $j < 0$, $a_0 > 0$, $a_0 + a_1 < 1$ and $\sum_{n=0}^{\infty} a_n = 1$. Let $\mu = \sum_{n=1}^{\infty} na_n$. It's known that $\{X_n\}$ is positive recurrent when $\mu < 1$; is null recurrent when $\mu = 1$; and is transient when $\mu > 1$. In this paper, we shall discuss the first returning speed and the last exit speed more precisely by means of $\{a_n\}$.

1. INTRODUCTION

Let $X = \{X_n : n = 0, 1, 2, \dots\}$ be a Markov chain with state space $\{0, 1, \dots\}$, transition matrix

$$P = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & \cdots \\ a_0 & a_1 & a_2 & a_3 & \cdots \\ 0 & a_0 & a_1 & a_2 & \cdots \\ 0 & 0 & a_0 & a_1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}.$$

That is, $p_{ij} = a_{j-(i-1)+}$, where $a_k = 0$ for all $k < 0$. Suppose that $0 < a_0 < 1$, $a_0 + a_1 < 1$ and $\sum_{n=0}^{\infty} a_n = 1$. This Markov chain is known as model repair shop, see [5] and [1]. The process X is an irreducible Markov chain. Let $\mu = \sum_{n=1}^{\infty} na_n$. Then X is positive recurrent if $\mu < 1$; is null recurrent if $\mu = 1$; and is transient if $\mu > 1$.

Suppose that $P(X_0 = 0) = 1$, $\tau = \tau_0 = \inf\{n \geq 1 : X_n = 0\}$ and $L = L_0 = \sup\{n \geq 0 : X_n = 0\}$. When $\mu = 1$, X is null recurrent. That is, $P(\tau < \infty) = 1$ and $E(\tau) = \infty$. But for what $0 < \alpha < 1$ that can make $E(\tau^\alpha) < \infty$? Similarly, when $\mu < 1$, X is positive recurrent. That is, $P(\tau < \infty) = 1$ and $E(\tau) < \infty$. But for what $1 < \alpha < \infty$ that can make $E(\tau^\alpha) < \infty$? In a word, when $\mu \leq 1$, the first returning time to 0 is finite almost surely, but how can we describe the returning speed in more detail? When $\mu > 1$, X is

Key words and phrases. repair shop, the first returning time, the last exit time, Markov chain.

*Department of mathematics, Hangzhou Normal University.

*Research supported by NSFC 11001070.

**Department of Mathematics, Zhejiang University.

**Research supported in part by NSFC 10601047 and by Zhejiang Provincial Natural Science Foundation of China grant J20091364.

**Corresponding author.

***Department of mathematics, Zhejiang University.

2000 AMS Classification: 60J10.

transient, that is $P(L < \infty) = 1$. On another words, the last exit time from 0 is finite almost surely. How can we characterize the exit speed more precisely? For example, for what $0 < \alpha < \infty$ that can make $E(L^\alpha) < \infty$? In this paper, we shall answer these questions.

This paper is organized as follows. In §2, the first returning speed and the last exit speed for any Markov chain are discussed. Then in the following, we focus on the repair shop model Markov chain. In §3, the first returning speed is discussed in the case that $\mu = 1$. In §4, the last exit speed is discussed in the case that $\mu > 1$. The first returning speed in the case that $\mu < 1$ is discussed in §5.

Throughout this paper, let $\mathbb{N}$ be the set of all positive integers. For any sequence $\{w(n)\}$, let $\Delta w(n) = w(n) - w(n-1)$ and $\Delta^2 w(n) = \Delta w(n) - \Delta w(n-1)$. Let $\mathcal{W}$ be the class of all nonnegative increasing sequence $(w(0), w(1), w(2), \dots)$ such that $\Delta w(n)$ is decreasing and $\lim_{n \rightarrow \infty} \Delta w(n) = 0$. For any two nonnegative sequences $\{a_n\}$ and $\{b_n\}$, we use $a_n \sim b_n$ to denote the case that there is $0 < c_1 < c_2 < \infty$ and $M \in \mathbb{N}$ such that $c_1 a_n \leq b_n \leq c_2 a_n$ for all $n \geq M$. For any nonnegative functions $f(x)$ and $g(x)$, $f(x) \sim g(x)$ as $x \rightarrow 0$ means that there is $0 < c_1 < c_2 < \infty$ and $0 < x_1 < 1$ such that $c_1 f(x) \leq g(x) \leq c_2 f(x)$ for all $0 < x < x_1$. Use $f(x) = o(g(x))$ as $x \rightarrow 0$ to denote the case that $\lim_{x \downarrow 0} \frac{f(x)}{g(x)} = 0$. If $\lim_{x \downarrow 0} \frac{f(x)}{g(x)} = \infty$, then we write $f(x) = \Theta(g(x))$ as $x \rightarrow 0$. Similarly $f(x) \sim g(x)$ as $x \rightarrow 1$ means that there is $0 < c_1 < c_2 < \infty$ and $0 < x_1 < 1$ such that $c_1 f(x) \leq g(x) \leq c_2 f(x)$ for all $x_1 < x < 1$.

2. GENERAL RESULTS FOR MARKOV CHAINS

Suppose that $\{X_n\}$ is a Markov chain on a probability space $(\Omega, \mathcal{F}, P)$, with n -step transition probabilities $p_{xy}^{(n)}$. Suppose that x is a state and $P(X_0 = x) = 1$. Let τ and L be the first returning time to x and the last exit time from x respectively. That is

$$\begin{aligned}\tau &= \inf\{n \geq 1 : X_n = x\}, \\ L &= \sup\{n \geq 0 : X_n = x\},\end{aligned}$$

where $\inf \emptyset = \infty$ and $\sup \emptyset = 0$. For any $t > 0$, let

$$\begin{aligned}F(t) &= E(t^\tau; \tau < \infty), \\ U(t) &= \sum_{n=0}^{\infty} p_{xx}^{(n)} t^n, \\ L(t) &= E(t^L; L < \infty).\end{aligned}$$

It is known that

$$(2.1) \quad U(t) = \begin{cases} \frac{1}{1-F(t)}, & \text{if } F(t) < 1; \\ \infty, & \text{otherwise.} \end{cases}$$

Let $p = F(1) = P(\tau < \infty)$, $q = 1 - p$. Then for any nonnegative integer n ,

$$P(L = n) = P(X_n = x, X_{n+1} \neq x, X_{n+2} \neq x, \dots) = qP(X_n = x).$$

Hence we have

$$(2.2) \quad L(t) = qU(t).$$

The state x is said to be recurrent if $p = 1$, otherwise x is said to be transient. If x is recurrent, then $P(L = \infty) = 1$, otherwise $P(L < \infty) = 1$. That is, if x is transient, then with probability one $\{X_n\}$ will leave from x sooner or later. Now a natural question has arisen: **how can we describe the last exit speed from x more precisely?** In [6], the moments of last exit times for Lévy processes are studied. In [8], the weighted moments of last exit times for Markov processes are studied. In this section, we shall use $F(t)$ and $U(t)$ to discuss the last exit speed for Markov chains.

On another hand, if x is recurrent, then though with probability one $\{X_n\}$ will return to x , but the first return speed may be quite different. To deal with this, recurrent states are divided into two classes: positive recurrent and null recurrent. The state x is positive recurrent if and only if $E(\tau) < \infty$. In [7], we discussed the first returning speed of null recurrent Markov chains. There, we proved that if x is a null recurrent state, then for any $w \in \mathcal{W}$,

$$E(w(\tau)) < \infty \text{ if and only if } \sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[F(1) - F(1 - \frac{1}{n})] < \infty.$$

But there are many positive recurrent Markov chains, and their returning speed may be quite different too. So another question has arisen: **how can we characterize the first returning speed of positive recurrent states more precisely?** In this section, we shall also discuss this question.

Let R_0 be the decay parameter of x , that is

$$R_0 = \sup\{t \geq 0 : F(t) \leq 1\}.$$

Then by equation 2.1, $R_0 = \sup\{t \geq 0 : U(t) < \infty\}$. If $F(R_0) = 1$, equivalently, $U(R_0) = \infty$, then we say $\{X_n\}$ is R_0 -recurrent. If $F'(R_0) < \infty$, that is, $E(R_0^\tau \tau; \tau < \infty) < \infty$, then x is said to be R_0 -positive recurrent. In this case, $\{X_n\}$ has some nice properties, see for example [3] and [4]. For discuss conveniently, we introduce another parameter

$$R_1 = \sup\{t \geq 0 : F(t) < \infty\}.$$

Obviously, $R_1 \geq R_0 \geq 1$. If X is recurrent, then $R_0 = 1$. If X is R_0 -transient or R_0 -null recurrent, then $R_1 = R_0$. The main results of this section are the following two theorems. Theorem 2.1 is followed by lemma 2.2 directly.

Theorem 2.1. *Suppose that $\{X_n\}$ is R_0 -recurrent, $R_1 < \infty$ and $F(R_1) < \infty$. Then we have the following properties.*

- (1) *For any $k \in \mathbb{N}$, $E(R_1^\tau \tau^k; \tau < \infty) < \infty$ if and only if $F^{(k)}(R_1) < \infty$.*
- (2) *If there is an integer $k \geq 0$ such that $F^{(k)}(R_1) < \infty$ but $F^{(k+1)}(R_1) = \infty$, then for any $w \in \mathcal{W}$, $E(R_1^\tau \tau^k w(\tau); \tau < \infty) < \infty$ if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[F^{(k)}(R_1) - F^{(k)}(R_1 - \frac{1}{n})] < \infty$.*

Theorem 2.2. *Suppose that $\{X_n\}$ is R_0 -transient. Then we have the following properties.*

(1) *That $R_1 = R_0$, $E(R_0^\tau; \tau < \infty) < 1$ and $E(R_0^L) < \infty$.*

(2) *For any $k \in \mathbb{N}$, the following four conditions are equivalent to each other.*

(a) $E(R_0^\tau \tau^k; \tau < \infty) < \infty$.

(b) $F^{(k)}(R_0) < \infty$.

(c) $U^{(k)}(R_0) < \infty$.

(d) $E(R_0^L L^k) < \infty$.

(3) *If there is an integer $k \geq 0$ such that $F^{(k)}(R_0) < \infty$ but $F^{(k+1)}(R_0) = \infty$, then for any $w \in \mathcal{W}$, the following four conditions are equivalent to each other.*

(i) $E(R_0^\tau \tau^k w(\tau); \tau < \infty) < \infty$.

(ii) $E(R_0^L L^k w(L)) < \infty$.

(iii) $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[F^{(k)}(R_0) - F^{(k)}(R_0 - \frac{1}{n})] < \infty$.

(iv) $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[U^{(k)}(R_0) - U^{(k)}(R_0 - \frac{1}{n})] < \infty$.

PROOF. We need only to show (2) and (3).

(2) It suffices to show that for any $k \in \mathbb{N}$, $F^{(k)}(R_0) < \infty$ if and only if $U^{(k)}(R_0) < \infty$. The equation $U(t) = 1 + U(t)F(t)$ implies that

$$(2.3) \quad U^{(k)}(t) = \sum_{i=0}^k \binom{k}{i} U^{(i)}(t) F^{(k-i)}(t), \forall 0 \leq t \leq R_0.$$

It follows that

$$(2.4) \quad U^{(k)}(R_0)(1 - F(R_0)) = U(R_0)F^{(k)}(R_0) + \sum_{i=1}^{k-1} \binom{k}{i} U^{(i)}(R_0)F^{(k-i)}(R_0).$$

Particularly, $U'(R_0)(1 - F(R_0)) = U(R_0)F'(R_0)$. Since $F(R_0) < 1$ and $U(R_0) < \infty$, $U'(R_0) < \infty$ if and only if $F'(R_0) < \infty$. Suppose that $k \geq 2$ and we have showed that for any $n < k$, $F^{(n)}(R_0) < \infty$ if and only if $U^{(n)}(R_0) < \infty$. By equation 2.4, $U^{(k)}(R_0) < \infty$ implies that $F^{(k)}(R_0) < \infty$. Conversely, if $F^{(k)}(R_0) < \infty$, then $F^{(n)}(R_0) < \infty$ for all $n < k$, and hence $U^{(n)}(R_0) < \infty$ for all $n < k$. Since $F(R_0) < 1$ and $U(R_0) < \infty$, by (2.4), $U^{(k)}(R_0) < \infty$. Thus our result holds by induction.

(3) By (2) and by Lemma 2.2, it suffices to show that for any integer $k \geq 0$,

$$(2.5) \quad U^{(k)}(R_0) - U^{(k)}(R_0 - x) \sim F^{(k)}(R_0) - F^{(k)}(R_0 - x), \text{ as } x \rightarrow 0.$$

If $k = 0$, then

$$U(R_0) - U(R_0 - x) = \frac{F(R_0) - F(R_0 - x)}{(1 - F(R_0))(1 - F(R_0 - x))} \sim F(R_0) - F(R_0 - x), \text{ as } x \rightarrow 0.$$

Now suppose that $k \geq 1$. By equation 2.4,

$$\begin{aligned} & U^{(k)}(R_0)(1 - F(R_0)) - U^{(k)}(R_0 - x)(1 - F(R_0 - x)) \\ = & [U(R_0)F^{(k)}(R_0) - U(R_0 - x)F^{(k)}(R_0 - x)] \\ & + \sum_{i=1}^{k-1} \binom{k}{i} [U^{(i)}(R_0)F^{(k-i)}(R_0) - U^{(i)}(R_0 - x)F^{(k-i)}(R_0 - x)]. \end{aligned}$$

Since $F'(R_0) < \infty$, $F(R_0) - F(R_0 - x) \sim x$ as $x \rightarrow 0$. Since $F^{(k+1)}(R_0) = \infty$, $U^{(k+1)}(R_0) = \infty$ and hence $x = o(U^{(k)}(R_0) - U^{(k)}(R_0 - x))$ as $x \rightarrow 0$. In addition $U^{(k)}(R_0) < \infty$. Thus

$$\begin{aligned} & U^{(k)}(R_0)(1 - F(R_0)) - U^{(k)}(R_0 - x)(1 - F(R_0 - x)) \\ = & [U^{(k)}(R_0) - U^{(k)}(R_0 - x)][1 - F(R_0)] + U^{(k)}(R_0 - x)[F(R_0 - x) - F(R_0)] \\ \sim & U^{(k)}(R_0) - U^{(k)}(R_0 - x), \text{ as } x \rightarrow 0. \end{aligned}$$

Similarly, as $x \rightarrow 0$,

$$\begin{aligned} & U(R_0)F^{(k)}(R_0) - U(R_0 - x)F^{(k)}(R_0 - x) \sim F^{(k)}(R_0) - F^{(k)}(R_0 - x), \\ & \sum_{i=1}^{k-1} \binom{k}{i} [U^{(i)}(R_0)F^{(k-i)}(R_0) - U^{(i)}(R_0 - x)F^{(k-i)}(R_0 - x)] \sim x. \end{aligned}$$

The fact that $F^{(k+1)}(R_0) = \infty$ yields that $x = o(F^{(k)}(R_0) - F^{(k)}(R_0 - x))$ as $x \rightarrow 0$. Therefore

$$(2.6) \quad U^{(k)}(R_0) - U^{(k)}(R_0 - x) \sim F^{(k)}(R_0) - F^{(k)}(R_0 - x), \text{ as } x \rightarrow 0.$$

□

Now we shall give two lemmas that are used in the proof of the theorems above. Suppose that $\{a_0, a_1, a_2, \dots\}$ is a nonnegative sequence. For any positive integer N , let $b_m^{(N)} = \sum_{n=mN}^{(m+1)N-1} a_n$ and $A_N = \sum_{n=0}^N a_n$. Let $A(t) = \sum_{n=0}^{\infty} a_n t^n$, $t \geq 0$. Let $h(n)$ be an increasing positive sequence such that

$$0 < \liminf_{n \rightarrow \infty} h(n)^n \leq \limsup_{n \rightarrow \infty} h(n)^n < 1.$$

The idea of Lemma 2.1 comes from Theorem 1 of [2].

Lemma 2.1. *If there is N_0 and $x > 0$ such that for all $N \geq N_0$,*

$$b_m^{(N)} \leq x b_0^{(N)}, \forall m.$$

Then $A_n \sim A(h(n))$.

PROOF. Obviously, $0 < h(n) < 1$ for all n . So we need only to consider the case that $\sum_{n=0}^{\infty} a_n = \infty$. For any N ,

$$\begin{aligned} A(h(N)) &= \sum_{n=0}^{\infty} a_n h(N)^n \geq \sum_{n=0}^N a_n h(N)^n \geq h(N)^N \sum_{n=0}^N a_n \\ &= A_N h(N)^N. \end{aligned}$$

On another hand, for any $N \geq N_0$,

$$\begin{aligned}
A(h(N)) &= \sum_{n=0}^{\infty} a_n h(N)^n = \sum_{m=0}^{\infty} \sum_{n=mN}^{(m+1)N-1} a_n h(N)^n \\
&\leq \sum_{m=0}^{\infty} h(N)^{mN} \sum_{n=mN}^{(m+1)N-1} a_n \leq \sum_{m=0}^{\infty} h(N)^{mN} (x \sum_{n=0}^N a_n) \\
&= \frac{x A_N}{1 - h(N)^N}.
\end{aligned}$$

Hence $A_n \sim A(h(n))$. □

By Lemma 2.1, if c is a positive constant and $\{a_n\}$ is a nonnegative decreasing sequence, then $A_n \sim A(1 - \frac{1}{cn})$.

Now suppose that $\sum_{n=0}^{\infty} a_n < \infty$. Then for any positive integer k and $0 \leq t < 1$,

$$A^{(k)}(t) = \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) a_n t^{n-k}.$$

Let

$$A^{(k)}(1) = \lim_{t \uparrow 1} A^{(k)}(t) = \sum_{n=k}^{\infty} a_n n(n-1) \cdots (n-k+1).$$

Then $\sum_{n=0}^{\infty} a_n n^k < \infty$ if and only if $A^{(k)}(1) < \infty$. Suppose that $\sum_{n=0}^{\infty} a_n n^k < \infty$ and $\sum_{n=0}^{\infty} a_n n^{k+1} = \infty$. Let $w \in \mathcal{W}$. Now we show consider the finiteness of $\sum_{n=0}^{\infty} a_n n^k w(n)$.

Lemma 2.2. *Suppose that c is a positive constant. Then $\sum_{n=0}^{\infty} a_n n^k w(n) < \infty$ if and only if $\sum_{n=1}^{\infty} -n \Delta^2 w(n+1) [A^{(k)}(1) - A^{(k)}(1 - \frac{1}{cn})] < \infty$.*

PROOF. We may assume that $w(0) = 0$ w.l.o.g. Let $b_m = \sum_{n=m}^{\infty} a_n n^k$. Then $b_m \leq b_0 < \infty$ and

$$\begin{aligned}
\sum_{n=0}^{\infty} a_n n^k w(n) &= \sum_{n=1}^{\infty} a_n n^k \sum_{m=1}^n \Delta w(m) = \sum_{m=1}^{\infty} \Delta w(m) \sum_{n=m}^{\infty} a_n n^k \\
&= \sum_{m=1}^{\infty} \Delta w(m) b_m = \sum_{m=1}^{\infty} b_m \sum_{n=m}^{\infty} [\Delta w(n) - \Delta w(n+1)] \\
&= \sum_{n=1}^{\infty} [\Delta w(n) - \Delta w(n+1)] \sum_{m=1}^n b_m.
\end{aligned}$$

Since b_m is decreasing, by Lemma 2.1, $\sum_{m=1}^n b_m \sim B(1 - \frac{1}{cn})$, where $B(t) = \sum_{n=1}^{\infty} b_n t^{n-1}$. On another hand,

$$\begin{aligned} \frac{A^{(k)}(1) - A^{(k)}(t)}{1-t} &= \sum_{n=k+1}^{\infty} a_n n(n-1) \cdots (n-k+1) (1+t+\cdots+t^{n-k-1}) \\ &= \sum_{i=0}^{\infty} t^i \sum_{n=i+k+1}^{\infty} a_n n(n-1) \cdots (n-k+1) \\ &\sim \sum_{i=0}^{\infty} b_{i+k+1} t^i = \frac{B(t)}{t^k} \left[1 - \frac{\sum_{j=1}^k b_j t^{j-1}}{B(t)} \right] \text{ as } t \rightarrow 1. \end{aligned}$$

Immediately,

$$B(1) = \sum_{m=1}^{\infty} b_m = \sum_{m=1}^{\infty} \sum_{n=m}^{\infty} a_n n^k = \sum_{n=1}^{\infty} a_n n^{k+1} = \infty.$$

Hence $\frac{A^{(k)}(1) - A^{(k)}(t)}{1-t} \sim B(t)$, as $t \rightarrow 1$. Therefore $\sum_{m=1}^n b_m \sim n[A^{(k)}(1) - A^{(k)}(1 - \frac{1}{cn})]$ which yields our result. $\square$

By Lemma 2.2, if there is $R > 0$ and nonnegative integer k such that $\sum_{n=0}^{\infty} a_n R^n n^k < \infty$ and $\sum_{n=0}^{\infty} a_n R^n n^{k+1} = \infty$, then for any $w \in \mathcal{W}$, $\sum_{n=0}^{\infty} a_n R^n n^k w(n) < \infty$ if and only if $\sum_{n=1}^{\infty} -n \Delta^2 w(n+1) [A^{(k)}(R) - A^{(k)}(R - \frac{1}{n})] < \infty$.

3. THE FIRST RETURNING SPEED FOR NULL RECURRENT REPAIR SHOP MODEL

In the following, suppose that $X = \{X_n : n = 0, 1, 2, \dots\}$ is a repair shop model Markov chain. That is, X is a Markov chain with state space $\{0, 1, \dots\}$, transition probability $p_{ij} = a_{j-(i-1)+}$, where $a_k = 0$ for all $k < 0$, $0 < a_0 < 1$, $a_0 + a_1 < 1$ and $\sum_{n=0}^{\infty} a_n = 1$.

Let $\mu = \sum_{n=1}^{\infty} n a_n$. Suppose that $P(X_0 = 0) = 1$ and $\tau = \inf\{n \geq 1 : X_n = 0\}$. For any $t > 0$, let $G(t) = \sum_{n=0}^{\infty} a_n t^n$ and $F(t) = E(t^\tau; \tau < \infty)$. Then by [5], for any $t > 0$,

$$(3.1) \quad F(t) = tG(F(t)).$$

In this section we shall consider the case that $\mu = 1$, that is, X is null recurrent.

Firstly, we shall characterize the asymptotic behavior of $1 - F(t)$ as $t \rightarrow 1$ by means of $G(t)$. Let $\mathcal{H}$ be the collection of all nonnegative strictly increasing continuous functions on $[0, 1]$. For any $f \in \mathcal{H}$, the inverse function of f exists, namely f^{-1} , defined on $[f(0), f(1)]$. For convenience, we may extend the domain of f^{-1} to $[f(0), \infty)$ by defining $f^{-1}(x) = 1$ for all $x \geq f(1)$. Then f^{-1} is strictly increasing on $[f(0), f(1)]$, and increasing on $[f(0), \infty)$.

For any $h \in [0, 1]$, let

$$\phi(h) = 1 - G'(1-h), \psi(h) = \int_0^h \phi(x) dx = G(1-h) - (1-h).$$

Since $\phi(h) > 0$ for all $0 < h < 1$, $\psi(h)$ is a strictly increasing continuous function on $[0, 1]$, that is $\psi \in \mathcal{H}$. Since $\psi(0) = 0$ and $\psi(1) = a_0$, ψ^{-1} is strictly increasing on $[0, a_0]$, increasing on $[0, \infty)$, $\psi^{-1}(0) = 0$ and $\psi^{-1}(x) = 1$ for all $x \geq a_0$.

Let $\mathcal{H}'$ be the collection of all $f \in \mathcal{H}$ that satisfying the following two properties:

(c1) For all $x, y \geq 0$ with $x + y \leq 1$, $f(x + y) \geq f(x) + f(y)$.

(c2) For any positive integer n and $x \in [0, \frac{1}{n}]$, $f(nx) \leq n^2 f(x)$.

For example, $f(x) = x^\alpha$ is in $\mathcal{H}'$ if and only if $1 \leq \alpha \leq 2$. We have the following lemmas.

Lemma 3.1. (1) That $\psi \in \mathcal{H}'$.

(2) For any $f \in \mathcal{H}'$, f^{-1} is a sub-additive function on $[0, \infty)$, that is for any $x, y \geq 0$,

$$f^{-1}(x + y) \leq f^{-1}(x) + f^{-1}(y).$$

(3) For any $f \in \mathcal{H}'$, any positive integer n and any $0 \leq y \leq f(\frac{1}{n})$,

$$f^{-1}(n^2 y) \geq n f^{-1}(y).$$

PROOF. (1) For any $h \in (0, 1)$, $\phi'(h) = G''(1 - h) > 0$, so ϕ is strictly increasing on $[0, 1]$.

Thus for any $h_1, h_2 \geq 0$ with $h_1 + h_2 \leq 1$,

$$\begin{aligned} \psi(h_1 + h_2) &= \int_0^{h_1} \phi(x) dx + \int_{h_1}^{h_1+h_2} \phi(x) dx \\ &\geq \int_0^{h_1} \phi(x) dx + \int_0^{h_2} \phi(x) dx = \psi(h_1) + \psi(h_2). \end{aligned}$$

Similarly, since $\phi''(h) = -G^{(3)}(1 - h) \leq 0$ on $(0, 1)$, $\phi'(h)$ is decreasing on $[0, 1]$ and hence $\phi(h)$ is a sub-additive function on $[0, 1]$. Thus for any $x \in [0, \frac{1}{n}]$,

$$\psi(nx) = \int_0^x n \phi(nh) dh \leq n^2 \int_0^x \phi(h) dh = n^2 \psi(x).$$

Consequently, $\psi \in \mathcal{H}'$.

(2) Suppose that $f \in \mathcal{H}'$. Then $f(0 + 0) \geq f(0) + f(0)$ implies that $f(0) = 0$. Hence the domain of f^{-1} is $[0, \infty)$, $f^{-1}(0) = 0$, $f^{-1}(x) = 1$ for all $x \geq f(1)$. For any $x, y \geq 0$, if $f^{-1}(x) + f^{-1}(y) \geq 1$, then $f^{-1}(x + y) \leq f^{-1}(x) + f^{-1}(y)$ holds.

If $f^{-1}(x) + f^{-1}(y) < 1$, then $f[f^{-1}(x) + f^{-1}(y)] \geq f[f^{-1}(x)] + f[f^{-1}(y)] = x + y$. It follows that $f^{-1}(x) + f^{-1}(y) \geq f^{-1}(x + y)$.

Therefore (2) holds.

(3) Let $x = f^{-1}(y)$. Then $0 \leq x \leq \frac{1}{n}$ and $f(nx) \leq n^2 f(x) = n^2 y$. Hence $f^{-1}(n^2 y) \geq n f^{-1}(y)$ holds. $\square$

Lemma 3.2. For any $0 < t < 1$, $1 - F(t) = \psi^{-1}(\frac{(1-t)F(t)}{t})$.

PROOF. For any $0 < t < 1$, by equation (3.1),

$$\psi(1 - F(t)) = G(F(t)) - F(t) = \frac{(1 - t)F(t)}{t}$$

which implies that $1 - F(t) = \psi^{-1}(\frac{(1-t)F(t)}{t})$. $\square$

Theorem 3.1. *As $t \rightarrow 1, 1 - F(t) \sim \psi^{-1}(1 - t)$.*

PROOF. Since $\lim_{t \rightarrow 1} \frac{F(t)}{t} = 1$, there is $0 < t_0 < 1$, such that for any $t_0 < t < 1$,

$$\frac{1}{2} \leq \frac{F(t)}{t} \leq 2.$$

Thus for any $t_0 < t < 1$,

$$\psi^{-1}\left[\frac{1}{2}(1 - t)\right] \leq \psi^{-1}\left[\frac{F(t)}{t}(1 - t)\right] \leq \psi^{-1}[2(1 - t)].$$

By Lemma 3.1 and Lemma 3.2, for any $t_0 < t < 1$,

$$\frac{1}{2}\psi^{-1}(1 - t) \leq 1 - F(t) \leq 2\psi^{-1}(1 - t).$$

Consequently, as $t \rightarrow 1, 1 - F(t) \sim \psi^{-1}(1 - t)$. $\square$

Particularly, $1 - F(1 - \frac{1}{n}) \sim \psi^{-1}(\frac{1}{n})$. By Theorem 2.1, we have the following corollary.

Corollary 3.1. (1) *For any $w \in \mathcal{W}$, $E[w(\tau)] < \infty$ if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n + 1)\psi^{-1}(\frac{1}{n}) < \infty$.*

(2) *For any $0 < \alpha < 1$, $E(\tau^\alpha) < \infty$ if and only if $\sum_{n=1}^{\infty} n^{\alpha-1}\psi^{-1}(\frac{1}{n}) < \infty$.*

Corollary 3.1 tells us that the returning speed of 0 can be characterized by ψ . The quicker that $\psi^{-1}(x)$ converges to 0 as x tends to 0, the quicker that $\{X_n\}$ returns to 0. Since ψ^{-1} is dependent on $G(x)$ completely, we can judge the finiteness of $E[w(\tau)]$ by means of $G(x)$ directly. Next, we shall discuss the returning speed in more detail.

Lemma 3.3. *Suppose that $\psi_1, \psi_2 \in \mathcal{H}'$ and n is a positive integer.*

(1) *If $\psi_1(x) \leq n\psi_2(x)$ as $x \rightarrow 0$, then $\psi_2^{-1}(y) \leq n\psi_1^{-1}(y)$ as $y \rightarrow 0$.*

(2) *If $\psi_1(x) \geq n^2\psi_2(x)$ as $x \rightarrow 0$, then $\psi_2^{-1}(y) \geq n\psi_1^{-1}(y)$ as $y \rightarrow 0$.*

PROOF. (1) There is $0 < x_0 < 1$ such that $\psi_1(x) \leq n\psi_2(x)$ for all $0 < x < x_0$. Let $y_0 = \psi_1(x_0)$. Then $y_0 > \psi_1(0) = 0$. For any $0 < y < y_0$, $0 < \psi_1^{-1}(y) < x_0$, and hence

$$\psi_1(\psi_1^{-1}(y)) \leq n\psi_2(\psi_1^{-1}(y)).$$

That is, $y \leq n\psi_2(\psi_1^{-1}(y))$. Consequently,

$$\psi_2^{-1}(y) \leq \psi_2^{-1}[n\psi_2(\psi_1^{-1}(y))] \leq n\psi_2^{-1}[\psi_2(\psi_1^{-1}(y))] = n\psi_1^{-1}(y).$$

(2) There is $0 < x_0 < \frac{1}{n}$ such that $\psi_1(x) \geq n^2\psi_2(x)$ for all $0 < x < x_0$. Thus by Lemma 3.1, for any $0 < x < x_0$,

$$\psi_2^{-1}[\psi_1(x)] \geq \psi_2^{-1}[n^2\psi_2(x)] \geq n\psi_2^{-1}[\psi_2(x)] = nx.$$

Letting $y = \psi_1(x)$, we get that $\psi_2^{-1}(y) \geq n\psi_1^{-1}(y)$ holds for all $0 < y < \psi_1(x_0)$. $\square$

By Lemma 3.3 and Theorem 3.1, we immediately get the following proposition.

Proposition 3.1. *Suppose that $\psi_1 \in \mathcal{H}'$.*

- (1) *If as $x \rightarrow 0$, $\psi(x) = o(\psi_1(x))$, then as $t \rightarrow 1$, $1 - F(t) = \Theta(\psi_1^{-1}(1 - t))$*
- (2) *If as $x \rightarrow 0$, $\psi(x) = \Theta(\psi_1(x))$, then as $t \rightarrow 1$, $1 - F(t) = o(\psi_1^{-1}(1 - t))$.*
- (3) *If as $x \rightarrow 0$, $\psi_1(x) \sim \psi(x)$, then as $t \rightarrow 1$, $1 - F(t) \sim \psi_1^{-1}(1 - t)$.*

Corollary 3.2. (1) *If $G''(1) < \infty$, then as $t \rightarrow 1$, $1 - F(t) \sim \sqrt{1 - t}$, and $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.*

(2) *If $G''(1) = \infty$, then as $t \rightarrow 1$, $1 - F(t) = o(\sqrt{1 - t})$, and $E(\tau^\alpha) < \infty$ for all $\alpha < \frac{1}{2}$.*

(3) *If there is $0 < \beta < 1$ such that $1 - G'(x) \sim (1 - x)^\beta$ as $x \rightarrow 1$, then as $t \rightarrow 1$, $1 - F(t) \sim (1 - t)^{\frac{1}{1+\beta}}$, and $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{\beta+1}$.*

PROOF. (1) Since $\lim_{x \uparrow 1} \frac{1 - G'(x)}{1 - x} = G''(1) < \infty$, as $h \rightarrow 0$, $\phi(h) = 1 - G'(1 - h) \sim h$. Thus as $x \rightarrow 0$,

$$\psi(x) = \int_0^x \phi(h) dh \sim \int_0^x h dh \sim x^2.$$

It is easy to verify that the function $\psi_1(x) := x^2$ is in $\mathcal{H}'$. By Proposition 3.1, as $t \rightarrow 1$, $1 - F(t) \sim \psi_1^{-1}(1 - t) = \sqrt{1 - t}$. By Theorem 2.1, $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

(2) Since $G''(1) = \infty$, as $h \rightarrow 0$, $\phi(h) = 1 - G'(1 - h) = \Theta(h)$. It implies that as $x \rightarrow 0$,

$$\psi(x) = \int_0^x \phi(h) dh = \Theta(x^2).$$

By Proposition 3.1 and Theorem 2.1, our result holds.

(3) As $h \rightarrow 0$, $\phi(h) = 1 - G'(1 - h) \sim h^\beta$. Thus as $x \rightarrow 0$,

$$\psi(x) = \int_0^x \phi(h) dh \sim \int_0^x h^\beta dh \sim x^{1+\beta}.$$

It is easy to verify that the function $\psi_2(x) = x^{1+\beta}$ is in $\mathcal{H}'$. By Proposition 3.1, as $t \rightarrow 1$, $1 - F(t) \sim \psi_2^{-1}(1 - t) = (1 - t)^{\frac{1}{1+\beta}}$. By Theorem 2.1, $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{1+\beta}$. $\square$

Proposition 3.1 and Corollary 3.2 tell us that the quicker that $G(x)$ tends to 1 as x tends to 1, the more slowly that $\{X_n\}$ returns to 0. In the case that $G''(1) < \infty$, $\{X_n\}$ returns to 0 the most slowly. In this case $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

Example 3.1. Suppose that $a_n = \frac{1}{2^{n+1}}$, $n \geq 0$. Then $\mu = 1$ and $\{X_n\}$ is null recurrent. It is easy to check that $G(t) = \frac{1}{2-t}$. Thus $F(t) = tG(F(t)) = \frac{t}{2-F(t)}$, which implies that $F(t) = 1 - \sqrt{1 - t}$. Particularly, $1 - F(1 - \frac{1}{n}) = \frac{1}{\sqrt{n}}$. Therefore $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

We may deduce this result by Corollary 3.2 directly. In fact $G'(t) = \frac{1}{(2-t)^2}$ and $G''(t) = \frac{2}{(2-t)^3}$. Hence $G''(1) = 2 < \infty$. So by Corollary 3.2, $1 - F(t) \sim \sqrt{1 - t}$, and $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

Example 3.2. Suppose that $a_0 = \frac{2}{3}$, $a_1 = 0$, $a_2 = \frac{1}{4}$, and for $n \geq 3$,

$$a_n = \frac{2}{3}(-1)^n \binom{\frac{3}{2}}{n} = \frac{1 \times 3 \times 5 \times \cdots \times (2n-5)}{2^{n-1}n!} > 0.$$

Then $G(x) = \sum_{i=0}^{\infty} a_i x^i = x + \frac{2}{3}(1-x)^{\frac{3}{2}}$ and $G'(x) = 1 - \sqrt{1-x}$. Thus $\sum_{n=0}^{\infty} a_n = G(1) = 1$ and $\mu = G'(1) = 1$. So $\{X_n\}$ is a null recurrent Markov chain. By $F(t) = t(G(F(t)))$, we get $\frac{(1-t)F(t)}{t} = \frac{2}{3}(1-F(t))^{\frac{3}{2}}$ which implies that $1-F(t) \sim (1-t)^{\frac{2}{3}}$. So $E(\tau^\alpha) < \infty$ if and only if $\alpha < \frac{2}{3}$.

We may also deduce this result by Corollary 3.2 directly. In fact, since $1 - G'(x) \sim (1-x)^{\frac{1}{2}}$ as $x \rightarrow 1$, by Corollary 3.2, $1 - F(t) \sim (1-t)^{\frac{2}{3}}$ as $t \rightarrow 1$.

4. THE LAST EXIT SPEED FOR TRANSIENT REPAIR SHOP MODEL

In this section, suppose that $\mu > 1$. Then $\{X_n\}$ is transient. We shall discuss the last exit speed from 0. Let $L = \sup\{n \geq 0 : X_n = 0\}$, R_0 be the decay parameter of $\{X_n\}$. If $F(R_0) = 1$, then $F(R_0) = R_0 G(F(R_0))$ implies that $R_0 = 1$. It contradicts that $F(1) < 1$. So $F(R_0) < 1$ and X is R_0 -transient. In addition, $F(t) = \infty$ provided $t > R_0$. For any $x \geq 0$ with $G'(x) < \infty$, let

$$\xi(x) = G(x) - xG'(x).$$

Then $\xi(0) = a_0 > 0$, $\lim_{t \uparrow 1} \xi(1) = 1 - \mu < 0$, and $\xi'(x) = -xG''(x) < 0$ for all $x > 0$. So there exists unique $0 < x_0 < 1$ such that $\xi(x_0) = 0$, that is, $G(x_0) = x_0 G'(x_0)$.

Theorem 4.1. (1) That $R_0 = \frac{x_0}{G(x_0)} > 1$, $F(R_0) = x_0 < 1$ and $E(R_0^L) < \infty$.

(2) That $E(R_0^\tau, \tau < \infty) = E(R_0^L) = \infty$.

(3) For any $w \in \mathcal{W}$, $E(R_0^\tau w(\tau); \tau < \infty) < \infty$ if and only if $E(R_0^L w(L)) < \infty$, and if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)\psi_{x_0}^{-1}(\frac{1}{n}) < \infty$, where $\psi_{x_0} = \frac{G(x_0(1-h))}{G(x_0)} - (1-h) \in \mathcal{H}'$ and $\psi_{x_0}^{-1}$ is the inverse function of ψ_{x_0} .

PROOF. (1) At first, we shall introduce a transformation. Suppose that $x > 0$ and $G(x) < \infty$. Define $P^{(x)}(X_0 = 0) := 1$. For any $n \geq 1$ and any nonnegative integers $i_0 = 0, i_1, \dots, i_n$, define

$$\begin{aligned} P^{(x)}(X_0 = 0, X_1 = i_1, \dots, X_n = i_n) &:= \frac{x^{\sum_{k=1}^n i_k - (i_{k-1}-1)^+}}{G(x)^n} P(X_0 = 0, X_1 = i_1, \dots, X_n = i_n) \\ &= \prod_{k=1}^n \frac{x^{i_k - (i_{k-1}-1)^+} a_{i_k - (i_{k-1}-1)^+}}{G(x)}. \end{aligned}$$

Let $\mathcal{F}_n = \sigma\{X_0, X_1, \dots, X_n\}$ and $\mathcal{F} = \sigma\{X_0, X_1, \dots\}$. It is easy to verify that $P^{(x)}$ is a pre-probability measure on $\bigcup_n \mathcal{F}_n$. Therefore $P^{(x)}$ has a unique extension to $\mathcal{F}$ which is denoted still by $P^{(x)}$. Then $P^{(x)}$ is a probability measure on $\mathcal{F}$. Under $P^{(x)}$, $\{X_n\}$ is a homogenous Markov chain with transition probability

$$p_{ij}^{(x)} = \frac{x^{j-(i-1)^+} a_{j-(i-1)^+}}{G(x)}.$$

Let

$$(4.1) \quad a_j^{(x)} = \frac{a_j x^j}{G(x)}, \forall j \geq 0.$$

Then $\sum_{j=0}^{\infty} a_j^{(x)} = \sum_{j=0}^{\infty} \frac{a_j x^j}{G(x)} = 1$, $a_0^{(x)} > 0$, $a_0^{(x)} + a_1^{(x)} = 1 - \sum_{j=2}^{\infty} a_j^{(x)} < 1$. Thus under $P^{(x)}$, $\{X_n\}$ is a repair shop model with $\{a_j^{(x)}\}$.

Next we shall calculate the value $F(t)$. For any $t > 0$, let

$$(4.2) \quad G^{(x)}(t) : = \sum_{j=0}^{\infty} a_j^{(x)} t^j = \frac{G(xt)}{G(x)},$$

$$(4.3) \quad F^{(x)}(t) : = E^{(x)}(t^\tau; \tau < \infty) = \sum_{n=1}^{\infty} P^{(x)}(\tau = n) t^n.$$

For any $n \geq 1$ and any nonnegative integers $i_0 = 0, i_1 \neq 0, i_2 \neq 0, \dots, i_{n-1} \neq 0, i_n = 0$,

$$P^{(x)}(X_0 = 0, X_1 = i_1, \dots, X_n = i_n) = \frac{x^{n-1}}{G(x)^n} P(X_0 = 0, X_1 = i_1, \dots, X_n = i_n).$$

It follows that

$$(4.4) \quad P^{(x)}(\tau = n) = \frac{x^{n-1}}{G(x)^n} P(\tau = n), \forall n$$

which implies that

$$F^{(x)}(t) = \frac{F(\frac{xt}{G(x)})}{x}.$$

Particularly, $F(\frac{x}{G(x)}) = xF^{(x)}(1) \leq x$.

Let $\mu^{(x)} = \sum_{j=1}^{\infty} j a_j^{(x)}$. Then

$$(4.5) \quad \mu^{(x)} = \frac{xG'(x)}{G(x)}.$$

If $\mu^{(x)} \leq 1$, then $\{X_n\}$ is recurrent under $P^{(x)}$, that is, $F^{(x)}(1) = 1$. Thus if $\frac{xG'(x)}{G(x)} \leq 1$, then $F(\frac{x}{G(x)}) = x$. For any $x \geq 0$ with $G(x) < \infty$, let

$$\eta(x) = \frac{x}{G(x)}.$$

Then $\eta'(x) = \frac{\xi(x)}{G(x)^2}$ for all x satisfying $G'(x) < \infty$. Thus $\eta'(x) > 0$ on $(0, x_0)$, and $\eta'(x) < 0$ when $x > x_0$. It leads to that $\eta(x)$ is strictly increasing on $[0, x_0]$, and then strictly decreasing when $x > x_0$. The maximum of $\eta(x)$ is $\eta(x_0)$. For any $y \geq 0$ with $F(y) < \infty$, $y = \frac{F(y)}{G(F(y))} = \eta(F(y)) \in [0, \eta(x_0)]$. That is, $F(y) < \infty$ if and only if $y \leq \frac{x_0}{G(x_0)}$. For any $y \in [0, \eta(x_0)]$, there is unique $x \in [0, x_0]$ such that $y = \eta(x)$. Since $\eta'(x) \geq 0$, $\frac{xG'(x)}{G(x)} \leq 1$ and $F(y) = x$, that is $F(y)$ is the minimum solution of $\eta(x) = y$.

By the discussion above, we get that $R_0 = \eta(x_0) = \frac{x_0}{G(x_0)}$ and $F(R_0) = x_0 < 1$. Since $x_0 \neq 1$, $R_0 = \eta(x_0) > \eta(1) = 1$. By Theorem 2.2, $E(R_0^L) < \infty$.

(2) By equation 4.5, $\mu^{(x_0)} = \frac{x_0 G'(x_0)}{G(x_0)} = 1$. It follows that $\{X_n\}$ is null recurrent under $P^{(x_0)}$. That is, $P^{(x_0)}(\tau < \infty) = 1$ and $E^{(x_0)}(\tau) = \infty$. By Corollary 3.1, for any $w \in \mathcal{W}$, $E^{(x_0)}(w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1) \psi_{x_0}^{-1}(\frac{1}{n}) < \infty$, where $\psi_{x_0}(h) = G^{(x_0)}(1-h) - (1-h)$. By equation 4.2, $\psi_{x_0}(h) = \frac{G(x_0(1-h))}{G(x_0)} - (1-h)$. By

equation 4.4, $E^{(x_0)}(\tau) = \frac{E(R_0^\tau; \tau < \infty)}{x_0}$ and $E^{(x_0)}(w(\tau)) = \frac{E(R_0^\tau w(\tau); \tau < \infty)}{x_0}$ for all $w \in \mathcal{W}$. Combination with Theorem 2.2, (2) and (3) hold. $\square$

Example 4.1. Suppose that $a_n = q^n p, n \geq 0$, where $0 < p < \frac{1}{2}$ and $q = 1 - p$. Then $G(t) = \frac{p}{1-tq}$ and $\mu = \frac{q}{p} > 1$. So $\{X_n\}$ is transient. By equation $F(t) = tG(F(t))$, we get that $F(t) = \frac{1-\sqrt{1-4pqt}}{2q}$. Thus $R_0 = \frac{1}{4pq}$, $F(R_0) = \frac{1}{2q} < 1$ and hence $\{X_n\}$ is R_0 -transient. Further $F'(R_0) = \infty$ and $F(R_0) - F(R_0 - \frac{1}{n}) = \frac{1}{2q} \sqrt{\frac{1}{nR_0}}$. Hence by Theorem 2.2, for any $\alpha > 0$, $E(R_0^L L^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

We may use Theorem 4.1 to deduce the results above. Clearly, $G'(t) = \frac{pq}{(1-tq)^2}$. Thus the equation $G(x) = xG'(x)$ has unique solution $x_0 = \frac{1}{2q}$. Hence $R_0 = \frac{x_0}{G(x_0)} = \frac{1}{4pq}$. In addition $a_n^{(x_0)} = \frac{a_n x_0^n}{G(x_0)} = \frac{1}{2^{n+1}}, n \geq 0$. That is under $P^{(x_0)}$, $\{X_n\}$ is the same process as in Example 3.1. Therefore for any $\alpha > 0$, $E(R_0^L L^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

5. THE FIRST RETURNING SPEED FOR POSITIVE RECURRENT REPAIR SHOP MODEL

In this section, suppose that $\mu < 1$. Then $\{X_n\}$ is positive recurrent. We shall discuss the returning time of 0. Let

$$R_1 = \sup\{t : F(t) < \infty\},$$

$$R = \sup\{t : G(t) < \infty\}.$$

Let $D = \{x \geq 0 : G(x) < \infty\}$ and $D' = \{x \geq 0 : G'(x) < \infty\}$. If $R = \infty$, then $D = D' = [0, \infty)$. If $R < \infty$ and $G'(R) < \infty$, then $G(R) < \infty$ and $D = D' = [0, R]$. If $R < \infty$, $G(R) < \infty$ but $G'(R) = \infty$, then $D = [0, R]$ and $D' = [0, R)$. If $R < \infty$ and $G(R) = \infty$, then $D = D' = [0, R)$.

Theorem 5.1. (1) If $G'(R) < \infty$ and $G(R) > RG'(R)$, then $R_1 = \frac{R}{G(R)}$ and $F(R_1) = R$.

(2) If $G'(R) = \infty$ or $G(R) \leq RG'(R)$, then there is unique $x_0 > 1$ such that $G(x_0) = x_0 G(x_0)$. In this case $R_1 = \frac{x_0}{G(x_0)} > 1$ and $F(R_1) = x_0 > 1$.

(3) That $R_1 = 1$ if and only if $R = 1$.

PROOF. For any $x \in D$, let

$$\eta(x) = \frac{x}{G(x)}.$$

For any $x \in D'$, let

$$\xi(x) = G(x) - xG'(x).$$

Then $\eta'(x) = \frac{\xi(x)}{G^2(x)}$ for all $x \in D'$. Since $\xi(0) = a_0 > 0$ and $\xi'(x) = -xG''(x) < 0$, $\xi(x) = 0$ has at most one solution. If $\xi(x) = 0$ has no solution, then $\xi > 0$ on D' and η is an increasing function on D . If $\xi(x) = 0$ has a unique solution namely x_0 , then $\xi > 0$ on $[0, x_0)$ and $\xi < 0$ on $(x_0, \infty) \cap D'$, which follows that η is strictly increasing on $[0, x_0]$ and then strictly decreasing when $x > x_0$. By the proof of Theorem 4.1, $F(y) < \infty$ if and only if $y = \eta(x)$ for some x with $G(x) < \infty$, and in this case,

$$F(y) = \{x : \eta(x) = y, G(x) \geq xG'(x)\} = \{x : \eta(x) = y, \eta'(x) \geq 0\} = \min\{x : \eta(x) = y\}.$$

If $\xi(x) = 0$ has a unique solution x_0 , then the range of $\eta(x)$ is $[0, \eta(x_0)]$. So $F(t) < \infty$ if and only if $t \leq \eta(x_0)$. That is $R_1 = \frac{x_0}{G(x_0)}$. Also $F(R_1) = x_0$. Since $\xi(1) = 1 - \mu > 0$, $x_0 > 1$ and $R_1 = \eta(x_0) > \eta(1) = 1$.

If $\xi(x) = 0$ has no solutions, then the range of $\eta(x)$ is $[0, \eta(R)]$ if $G(R) < \infty$, or $[0, \eta(R))$ if $G(R) = \infty$. We shall show that $G'(R) < 1$. If $R = 1$, then $G'(1) = \mu < 1$ holds. Now suppose that $R > 1$. Note that $\xi(x) > 0$ for all $x < R$ and $\eta(x)$ is strictly increasing on $[0, R]$. For any $\frac{1+R}{2} < x < R$,

$$G'(x) < \frac{G(x)}{x} = \frac{1}{\eta(x)} \leq \frac{1}{\eta(\frac{1+R}{2})}.$$

Thus

$$G'(R) = \lim_{x \uparrow R} G'(x) \leq \frac{1}{\eta(\frac{1+R}{2})} < \frac{1}{\eta(1)} = 1.$$

Therefore $G'(R) < 1$ and $G(R) < \infty$. Consequently, $R_1 = \eta(R) = \frac{R}{G(R)}$ and $F(R_1) = R$. If $R = 1$, then $R_1 = \eta(1) = 1$. If $R > 1$, then $R_1 = \eta(R) > \eta(1) = 1$. $\square$

Similar as the proof of the statements (2) and (3) in Theorem 4.1, we may prove the following theorem.

Theorem 5.2. *Suppose that there is unique $x_0 > 1$ such that $G(x_0) = x_0 G'(x_0)$. Then $E(R_1^\tau \tau) = \infty$, and for any $w \in \mathcal{W}$, $E(R_1^\tau w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} -n \Delta^2 w(n+1) \psi_{x_0}^{-1}(\frac{1}{n}) < \infty$, where $\psi_{x_0} = \frac{G(x_0(1-h))}{G(x_0)} - (1-h) \in \mathcal{H}'$ and $\psi_{x_0}^{-1}$ is the inverse function of ψ_{x_0} .*

We shall discuss the first returning speed of $\{X_n\}$ further in the case that $G(x) = xG'(x)$ has no positive solutions.

Lemma 5.1. *For any integer $n \geq 2$, $E(\tau^n) < \infty$ if and only if $G^{(n)}(1) < \infty$.*

PROOF. By $F(t) = tG(F(t))$, we get that

$$(5.1) \quad F'(t) = G(F(t)) + tG'(F(t))F'(t),$$

$$(5.2) \quad F''(t) = 2G'(F(t))F'(t) + tG''(F(t))[F'(t)]^2 + tG'[F(t)]F''(t).$$

We shall use induction to show that

$$(5.3) \quad F^{(n)}(t) = H_{n-1}(t) + tG^{(n)}[F(t)][F'(t)]^n + tG'[F(t)]F^{(n)}(t), \forall n \geq 2,$$

where $H_{n-1}(t)$ is a polynomial function of $F'(t), \dots, F^{(n-1)}(t), G'(F(t)), \dots, G^{(n-1)}(F(t))$ with positive coefficients.

By equation 5.2, equation 5.3 holds for $n = 2$. Suppose that equation 5.3 holds for some $n = k \geq 2$, that is

$$F^{(k)}(t) = H_{k-1}(t) + tG^{(k)}[F(t)][F'(t)]^k + tG'[F(t)]F^{(k)}(t).$$

Then

$$F^{(k+1)}(t) = H_k(t) + tG^{(k+1)}[F(t)][F'(t)]^{k+1} + tG'[F(t)]F^{(k+1)}(t),$$

where

$$\begin{aligned} H_k(t) &= H'_{k-1}(t) + G^{(k)}[F(t)][F'(t)]^k + ktG^{(k)}[F(t)][F'(t)]^{k-1}F''(t) \\ &\quad + G'[F(t)]F^{(k)}(t) + tG''[F(t)]F'(t)F^{(k)}(t). \end{aligned}$$

Obviously, $H_k(t)$ is a polynomial function of $F'(t), \dots, F^{(k)}(t), G'(F(t)), \dots, G^{(k)}(F(t))$ with positive coefficients. Thus equation 5.3 holds.

It follows that

$$(5.4) \quad F^{(n)}(t) = \frac{H_{n-1}(t) + tG^{(n)}[F(t)][F'(t)]^n}{1 - tG'[F(t)]}, \forall n \geq 2.$$

Particularly, by equation 5.1 and 5.4,

$$(5.5) \quad F'(1) = \frac{1}{1 - \mu},$$

$$(5.6) \quad F^{(n)}(1) = \frac{H_{n-1}(1) + G^{(n)}(1)\left(\frac{1}{1-\mu}\right)^n}{1 - \mu}, \forall n \geq 2.$$

Obviously, $F''(1) < \infty$ if and only if $G''(1) < \infty$. Suppose that when $n \leq k$, $F^{(n)}(1) < \infty$ if and only if $G^{(n)}(1) < \infty$.

If $F^{(k+1)}(1) < \infty$, then $F^{(n)}(1) < \infty$ for all $n \leq k$, and hence $G^{(n)}(1) < \infty$ for all $n \leq k$, which yields that $H_k(1) < \infty$. Consequently $G^{(k+1)}(1) < \infty$. Similarly, if $G^{(k+1)}(1) < \infty$, then $F^{(k+1)}(1) < \infty$.

Therefore by induction, for all $n \geq 2$, $F^{(n)}(1) < \infty$ if and only if $G^{(n)}(1) < \infty$. Now the fact that $E(\tau^n) < \infty$ if and only if $F^{(n)}(1) < \infty$ yields our result. $\square$

Lemma 5.2. *Suppose that there is $k \in \mathbb{N}$ such that $E(\tau^k) < \infty$ but $E(\tau^{k+1}) = \infty$. Then for any $w \in \mathcal{W}$, $E(\tau^k w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} n^k w(n) a_n < \infty$.*

PROOF. By Lemma 5.1, $G^{(k)}(1) < \infty$ but $G^{(k+1)}(1) = \infty$. By Theorem 2.1, for any $w \in \mathcal{W}$, $E(\tau^k w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[F^{(k)}(1) - F^{(k)}(1 - \frac{1}{n})] < \infty$. By Lemma 2.2, for any $w \in \mathcal{W}$, $\sum_{n=1}^{\infty} n^k w(n) a_n < \infty$ if and only if $\sum_{n=1}^{\infty} -n\Delta^2 w(n+1)[G^{(k)}(1) - G^{(k)}(1 - \frac{1}{n})] < \infty$.

Thus it suffices to show that

$$(5.7) \quad F^{(k)}(1) - F^{(k)}(1-t) \sim G^{(k)}(1) - G^{(k)}(1-t), \text{ as } t \rightarrow 0.$$

Firstly, we shall show that

$$(5.8) \quad G^{(k)}(1) - G^{(k)}[F(1-t)] \sim G^{(k)}(1) - G^{(k)}(1-t), \text{ as } t \rightarrow 0.$$

For $0 < t < 1$, $F(1-t) \leq 1-t$ and hence $G^{(k)}(1) - G^{(k)}[F(1-t)] \geq G^{(k)}(1) - G^{(k)}(1-t)$. Since the function $G^{(k)}(x)$ is convex,

$$G^{(k)}(1) - G^{(k)}[F(1-t)] \leq [1 - F(1-t)] \frac{G^{(k)}(1) - G^{(k)}(1-t)}{t} \leq F'(1)[G^{(k)}(1) - G^{(k)}(1-t)].$$

Consequently, 5.8 holds.

If $k = 1$, then

$$F'(1) - F'(1-t) = \frac{1 - (1-t)G'[F(1-t)] - (1-\mu)G[F(1-t)]}{(1-\mu)\{1 - (1-t)G'[F(1-t)]\}}.$$

Obviously,

$$\begin{aligned} & 1 - (1-t)G'[F(1-t)] - (1-\mu)G[F(1-t)] \\ = & (1-\mu)\{G[F(1)] - G[F(1-t)]\} + \{\mu - G'[F(1-t)]\} + tG'[F(1-t)]. \end{aligned}$$

Note that $G'(1) < \infty$ and $F'(1) < \infty$, combination with 5.8, we get that as $t \rightarrow 0$,

$$\begin{aligned} (1-\mu)\{G[F(1)] - G[F(1-t)]\} & \sim t, \\ \mu - G'[F(1-t)] & \sim G'(1) - G'(1-t) \\ tG'[F(1-t)] & \sim t, \end{aligned}$$

and

$$(1-\mu)\{1 - (1-t)G'[F(1-t)]\} \sim (1-\mu)^2.$$

The fact that $G''(1) = \infty$ implies that $t = o(G'(1) - G'(1-t))$ as $t \rightarrow 0$. Therefore as $t \rightarrow 0$, $F'(1) - F'(1-t) \sim G'(1) - G'(1-t)$.

If $k \geq 2$, then by equation 5.3,

$$\begin{aligned} F^{(k)}(1) - F^{(k)}(1-t) & = \frac{H_{k-1}(1) - H_{k-1}(1-t)}{1 - (1-t)G'[F(1-t)]} \\ & + \frac{\{\mu - (1-t)G'[F(1-t)]\}[H_{k-1}(1) + G^{(k)}(1)(\frac{1}{1-\mu})^k]}{(1-\mu)\{1 - (1-t)G'[F(1-t)]\}} \\ & + \frac{G^{(k)}(1)(\frac{1}{1-\mu})^k - (1-t)G^{(k)}[F(1-t)][F'(1-t)]^k}{1 - (1-t)G'[F(1-t)]} \end{aligned}$$

By 5.8 and by the fact that $t \sim o(G^{(k)}(1) - G^{(k)}(1-t))$ as $t \rightarrow 0$, we get

$$\begin{aligned} & G^{(k)}(1)(\frac{1}{1-\mu})^k - (1-t)G^{(k)}[F(1-t)][F'(1-t)]^k \\ = & (1-t)(\frac{1}{1-\mu})^k \{G^{(k)}(1) - G^{(k)}[F(1-t)]\} + (1-t)G^{(k)}[F(1-t)]\{(\frac{1}{1-\mu})^k - [F'(1-t)]^k\} \\ & + tG^{(k)}(1)(\frac{1}{1-\mu})^k \\ \sim & G^{(k)}(1) - G^{(k)}(1-t), \text{ as } t \rightarrow 0. \end{aligned}$$

In addition, as $t \rightarrow 0$,

$$\begin{aligned} & H_{k-1}(1) - H_{k-1}(1-t) \sim t, \\ & \{\mu - (1-t)G'[F(1-t)]\}[H_{k-1}(1) + G^{(k)}(1)(\frac{1}{1-\mu})^k] \sim t, \\ & 1 - (1-t)G'[F(1-t)] \sim 1 - \mu. \end{aligned}$$

Therefore

$$F^{(k)}(1) - F^{(k)}(1-t) \sim G^{(k)}(1) - G^{(k)}(1-t) \text{ as } t \rightarrow 0,$$

and our result holds. $\square$

Theorem 5.3. *If $G'(R) < \infty$ and $G(R) > RG'(R)$, then we have the following properties.*

- (1) *That $E(R_1^\tau \tau) < \infty$,*
- (2) *For any $\alpha > 1$, $E(R_1^\tau \tau^\alpha) < \infty$ if and only if $\sum_{n=1}^{\infty} R^n n^\alpha a_n < \infty$.*
- (3) *If there is $k \in \mathbb{N}$ such that $E(R_1^\tau \tau^k) < \infty$ but $E(R_1^\tau \tau^{k+1}) = \infty$, then for any $w \in \mathcal{W}$, $E(R_1^\tau \tau^k w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} R^n n^k w(n) a_n < \infty$.*

PROOF. Since $\mu^{(R)} = \frac{RG'(R)}{G(R)} < 1$, $\{X_n\}$ is still positive recurrent under $P^{(R)}$. Hence $E^{(R)}(\tau) < \infty$. By Lemma 5.1, for any integer $k \geq 2$, $E^{(R)}(\tau^k) < \infty$ if and only if $\sum_{n=1}^{\infty} a_n^{(R)} n^k < \infty$. In addition, by Lemma 5.2, if there is $k \in \mathbb{N}$ such that $E^{(R)}(\tau^k) < \infty$ but $E^{(R)}(\tau^{k+1}) = \infty$, then for any $w \in \mathcal{W}$, $E^{(R)}(\tau^k w(\tau)) < \infty$ if and only if $\sum_{n=1}^{\infty} n^k w(n) a_n^{(R)} < \infty$. On another hand, $E^{(R)}(\tau^k) = \frac{E(R_1^\tau \tau^k)}{R}$, $E^{(R)}(\tau^k w(\tau)) = \frac{E(R_1^\tau \tau^k w(\tau))}{R}$ and $a_n^{(R)} = \frac{a_n R^n}{G(R)}$. Thus our result holds. $\square$

Example 5.1. Suppose that $a_n = q^n p$, $n \geq 0$, where $\frac{1}{2} < p < 1$ and $q = 1 - p$. Then $G(t) = \frac{p}{1-tq}$ and $\mu = \frac{q}{p} < 1$. So $\{X_n\}$ is positive recurrent. Similar as Example 4.1, $F(t) = \frac{1-\sqrt{1-4pqt}}{2q}$. Thus $R_1 = \frac{1}{4pq}$, $F'(R_1) = \infty$ and $F(R_1) - F(R_1 - \frac{1}{n}) = \frac{1}{2q} \sqrt{\frac{1}{R_1 n}}$. So by Theorem 2.1, $E(R_1^\tau \tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

We may use Theorem 5.1 and Theorem 5.2 to deduce our result. Immediately, $R = \sup\{t : G(t) < \infty\} = \frac{1}{q}$ and $G'(R) = \infty$. Thus the equation $G(x) = xG'(x)$ has unique solution, indeed, it is $x_0 = \frac{1}{2q}$. Thus $R_1 = \frac{x_0}{G(x_0)} = \frac{1}{4pq}$. Also under $P^{(x_0)}$, $\{X_n\}$ is the same process as in Example 3.1. Therefore for any $\alpha > 0$, $E(R_1^\tau \tau^\alpha) < \infty$ if and only if $\alpha < \frac{1}{2}$.

Example 5.2. suppose that $\alpha > 2$ and $a_k = (\frac{1}{k+1})^\alpha - (\frac{1}{k+2})^\alpha$, $k \geq 0$. Then

$$\sum_{k=0}^{\infty} k a_k = (\frac{1}{2})^\alpha + (\frac{1}{3})^\alpha + \cdots \leq \int_1^{\infty} (\frac{1}{x})^\alpha dx = \frac{1}{\alpha-1} < 1.$$

Thus $\{X_n\}$ is a positive recurrent. As $k \rightarrow \infty$, $a_k \sim k^{-\alpha-1}$. So $G(t) = \infty$ for all $t > 1$ and $R = 1$. By Theorem 5.1, $R_1 = 1$. For any $\beta > 0$, $\sum_{k=1}^{\infty} a_k k^\beta < \infty$ if and only if $\beta < \alpha$. So by Theorem 5.3, $E(\tau^\beta) < \infty$ if and only if $\beta < \alpha$.

REFERENCES

- [1] Pierre Bremaud, MARKOV CHAINS: GIBBS FIELDS, MONTE CARLO SIMULATION, AND QUEUES, Springer-Verlag New York(1999).
- [2] Dean Isaacson and Peter Colwell, *Levels of Null Persistency for Markov Chains*, J.Appl.Prob, 19,425-429(1982).
- [3] D. Vere-Jones, *Geometric ergodicity in denumerable Markov Chains*, Quart. J. Math. Vol., 13, No. 2, 7-28(1962).
- [4] J. F. C. Kingman, *The exponential decay of Markov transition probabilities*, Proc. London Mth. Soc., Vol. 13, No.3, 337-358(1963).
- [5] S. Karlin and H. M. Taylor, A FIRST COURSE IN STOCHASTIC PROCESSES, Academic Press(1975).
- [6] Ken-iti Sato, and Toshiro Watanabe, *Moments of last exit times for Levy processes*, Ann. Inst. Henri Poincaré, Probab. Statist., Volume 40, Issue 2, 207-225(2004).

- [7] M. Z. Zhao, *The first returning speed of null recurrent Markov chains*, Chinese Ann. Math. Ser. A, Vol. 27, No. 6, 761-770(2006).
- [8] M. Z. Zhao and H. Z. Zhang, *The weighted transience and recurrence of Markov processes*, Acta Mathematica Sinica, English Series, Vol. 23, No. 1, 111-126(2007).